


BD SurePath[™] Liquid-based Pap Test Collection Procedure

Important: See specific instructions for each device type



1. Collect

Collect the cytology sample using either a broom-like device or combination brush/plastic spatula with detachable heads. Follow manufacturer's collection instructions for detachable-head device(s).



2. Drop

After preparing the conventional smear, use the interior rim of the BD SurePath[™] Collection Vial to pull off the head of the broom-like device or snap the heads of the brush and spatula into the large opening in the collection vial.



3. Send

Place the cap on the vial and tighten. Send the BD SurePath™ Collection Vial to the lab for processing.

Device Type

Directions For Use

Rovers[®] Cervex-Brush[®]



Insert the Rovers® Cervex-Brush® into the endocervical canal and rotate **5** times in a **clockwise** direction.

Rovers[®] Cervex-Brush[®] Combi



Insert the Rovers® Cervex-Brush® Combi into the endocervical canal and rotate **2** times in a **clockwise** direction.



Cytobrush[®] Plus GT



Insert Cytobrush® Plus GT into the endocervix until only the bottom-most bristles are exposed at the os. Slowly rotate 1/4 to 1/2 turn in one direction. To reduce unnecessary bleeding, do not over rotate brush.





Insert the contoured end of the Pap-Perfect[®] plastic spatula and rotate **360°** around the entire ectocervix.

えん

Contraindications:		
Rovers [®] Cervex-Brush [®]	As indicated in Rovers [®] Cervex-Brush [®] product label, the broom may be used on pregnant patients up to 10 weeks gestation.	
Rovers [®] Cervex-Brush [®] Combi	The Cervex-Brush® Combi should not be used on patients during pregnancy.	
Cytobrush® Plus GT	Do not use the Cytobrush® Plus GT on pregnant patients or for endometrial sampling.	

Becton Dickinson Pty Ltd 4 Research Park Drive, Macquarie University Research Park, North Ryde NSW 2113, Toll Free: 1800 656 100 BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2015. ANZDS134

